

bridgebio

hope through
rigorous science

Corporate presentation

November 2020



Forward-Looking Statements and Disclaimer

Statements in this Presentation that are not statements of historical fact are forward-looking statements within the meaning of Section 27A of the Securities Act of 1933, as amended, and Section 21E of the Securities Exchange Act of 1934, as amended. Such forward-looking statements include, without limitation, statements regarding BridgeBio Pharma, Inc.'s (the "Company's") research and clinical development plans, expected manufacturing capabilities, strategy, regulatory matters, market size and opportunity, future financial position, future revenue, projected costs, prospects, plans, objectives of management, and the Company's ability to complete certain milestones. Words such as "believe," "anticipate," "plan," "expect," "intend," "will," "may," "goal," "potential," "should," "could," "aim," "estimate," "predict," "continue" and similar expressions or the negative of these terms or other comparable terminology are intended to identify forward-looking statements, though not all forward-looking statements necessarily contain these identifying words. These forward-looking statements are neither forecasts, promises nor guarantees, and are based on the beliefs of the Company's management as well as assumptions made by and information currently available to the Company. Such statements reflect the current views of the Company with respect to future events and are subject to known and unknown risks, including business, regulatory, economic and competitive risks, uncertainties, contingencies and assumptions about the Company, including, without limitation, risks inherent in developing therapeutic products, the success, cost, and timing of the Company's product candidate development activities and ongoing and planned preclinical studies and clinical trials, trends in the industry, the legal and regulatory framework for the industry, the Company's ability to obtain and maintain regulatory approval for its product candidates, the Company's ability to commercialize its product candidates, future agreements with third parties in connection with the development or commercialization of the Company's product candidates, the size and growth potential of the market for the Company's product candidates, the accuracy of the Company's estimates regarding expenses, future revenue, future expenditures and needs for and ability to obtain additional financing, the Company's ability to obtain and maintain intellectual property protection for its product candidates, potential adverse impacts due to the global COVID-19 pandemic such as delays in clinical trials, preclinical work, overall operations, regulatory review, manufacturing and supply chain interruptions, adverse effects on healthcare systems and disruption of the global economy, and those risks and uncertainties described under the heading "Risk Factors" in the Company's most recent Quarterly Report on Form 10-Q and Annual Report on Form 10-K filed with the U.S. Securities and Exchange Commission ("SEC") and in subsequent filings made by the Company with the SEC, which are available on the SEC's website at www.sec.gov. In light of these risks and uncertainties, many of which are beyond the Company's control, the events or circumstances referred to in the forward-looking statements, expressly or implicitly, may not occur. The actual results may vary from the anticipated results and the variations may be material. You are cautioned not to place undue reliance on these forward-looking statements, which speak the Company's current beliefs and expectations only as of the date this Presentation is given. Except as required by law, the Company disclaims any intention or responsibility for updating or revising any forward-looking statements contained in this Presentation in the event of new information, future developments or otherwise. No representation is made as to the safety or effectiveness of these product candidates for the therapeutic use for which such product candidates are being studied.

Certain information contained in this Presentation relates to or is based on studies, publications, surveys and other data obtained from third-party sources and the Company's own internal estimates and research. While the Company believes these third-party sources to be reliable as of the date of this Presentation, it has not independently verified, and makes no representation as to the adequacy, fairness, accuracy or completeness of, any information obtained from third-party sources. In addition, all of the market data included in this Presentation involves a number of assumptions and limitations, and there can be no guarantee as to the accuracy or reliability of such assumptions. Finally, while the Company believes its own internal research is reliable, such research has not been verified by any independent source.

The Company is the owner of various trademarks, trade names and service marks. Certain other trademarks, trade names and service marks appearing in this Presentation are the property of third parties. Solely for convenience, the trademarks and trade names in this Presentation are referred to without the ® and TM symbols, but such references should not be construed as any indicator that their respective owners will not assert, to the fullest extent under applicable law, their rights thereto.

BridgeBio Pharma: Hope through rigorous science

Our mission: To **discover**, **create**, **test** and **deliver** transformative medicines to treat patients who suffer from genetic diseases and cancers with clear genetic drivers




BridgeBio corporate overview



Strategy



Platform




Products



We are at Day 1 in the era of genetic medicine

Hundreds of monogenic disease-causing variants are discovered every year...


...and common disease associations are increasing exponentially



A vast opportunity to help patients



Americans are living with a genetic diseases




Of people affected are children




Of these diseases have an approved therapy

US prevalence, mn



Source: Global Genes, American Cancer Society, American Heart Association, Alzheimer’s Association, Arthritis Foundation, Bailey *et al.*, *Cell* 2018

Our strategy is simple



History teaches us about strategy:

1. Right playing field



2. Right tenets



3. Stay adaptive




BBIO applications:

1. Genetic disease

2. Beautiful science, NPV positive

3. No initial focus on TA, disease, or modality. Repeated application

Our organizational principles enable scale



History teaches us about growth:

1. Simple rules repeated at many levels



2. De-centralized cities grow with *returns to scale*, centralized companies slow with economies of scale




BBIO applications:

1. Simple rules – put patients first, think independently and let science speak, be efficient

2. De-centralized approach – small teams that focus and are incented at the level of each asset, scale that allows for rapid failure, learning


Result: Pipeline momentum



Increasing returns to scale – BridgeBio since IPO

Achievement:

- NDA for MoCD Type A accepted, ODD & Fast Track received for 2L CCA program
- 7 INDs filed
- Twelve new clinical trials initiated (18 total), >400 trial sites across 26 countries
- 8 new programs, including LGMD2i and ADH1, both in the clinic
- TTR clinical data, DEB clinical data, CAH and Canavan pre-clinical data, achon pre-clinical data, TIO data



BridgeBio corporate overview



Strategy



Platform



Products



BridgeBio drug engineering basics: our platform

Discover


Novel genetic
disease targets



Well described diseases
than can be targeted at
their source

Create


Medicines with industry-
leading research capabilities



Tailored therapeutic
technologies to create first
or best-in-class medicines

Test

Our drugs through global
development footprint



Broad clinical development
capabilities across therapeutic
areas and geographies

Deliver

Our products to patients through
commercial infrastructure



Building the capabilities to
deliver genetic medicines to
patients globally



Discover

Capabilities to identify new genetic disease targets at scale

Our target identification engine is driven by three core areas of strength:

Computational genomics / statistical genetics

- Mining of large genotype-phenotype databases
- *De novo* target discovery
- Target validation
- Indication expansion

Systematic disease mapping

- Manual annotation and prioritization of the 7K known genetic diseases

Partnering with top academic researchers

- >15 current partnerships





Discover

Scientific insight and judgment from industry leaders with a proven track record



Charles Homcy, MD

Founder and Chairman of
Pharmaceuticals




Frank McCormick, PhD

Founder and Chairman of
Oncology



Richard Scheller, PhD

Chairman of R&D



Len Post, PhD

Advisor



Phil Reilly, MD, JD

Advisor





Create


We select the optimal therapeutic modality to target each disease at its source

Industry-leading capabilities across 4 modalities:

Medicinal chemistry

- Molecular dynamics
- Reversible and irreversible chemistry
- Topical formulations


Optimal use: Inhibition of GOF or allosteric activation of LOF mutations



Therapeutic proteins

- Large protein manufacturing
- Formulation expertise
- Comparability assay development

Optimal use: Replacement of extracellular protein in LOF diseases



Gene therapy

- Vector optimization
- Novel capsid engineering
- Analytical assay development


Optimal use: Replacement of intracellular protein in LOF diseases



Antisense oligonucleotides

- Target mapping with functional genomics
- Activity screening assay development
- Novel backbone and base chemistry

Optimal use: Inhibition of GOF or activation of WT allele in LOF diseases





Create

Research leaders with a productive history developing novel therapeutics

Mendelian



Uma Sinha, PhD
Chief Scientific Officer



Oncology



Robert Zamboni, PhD
Chemistry



Eli Wallace, PhD
Chief Scientific Officer, Oncology



Pedro Beltran, PhD
SVP, Oncology



Gene therapy



Clayton Beard, PhD
SVP, Research and Development






Our global clinical development footprint

Test

- **18 ongoing trials** across 5 different therapeutic areas, >400 trial sites, and 26 countries
- **Creative clinical and regulatory strategy**, e.g., unique, nested Phase 3 trial design for acoramadis in ATTR
- **Central operations toolkit** for enrollment, protocol quality, site activation, CRO quality, regional performance
- **Expert, dedicated R&D teams** in each therapeutic area
 - **Cardio/renal:** Jonathan Fox, MD, PhD  MyoKardia  AstraZeneca
 - **Oncology:** Susan Moran, MD  Puma Biotechnology  genzyme
 - **Gene Therapy:** Adam Shaywitz, MD, PhD  BIOMARIN  AMGEN





Deliver

Building capabilities to deliver our products to patients across the globe

- **Global commercial infrastructure** to leverage our drug and disease expertise
- **Diagnostic partnerships** to identify patients in need of our medicines
- **Disease awareness strategies** including close partnerships with patient advocacy groups
- **Country-specific Early Access Programs (EAP)** and patient assistance programs
- **Commercial partners in strategic geographies:**

*FGFRi and SHP2i
in China:*



*TTR
in Japan:*



*MoCD type A
in Israel:*



Key people: Matt Outten (CCO), Jennifer Cook (BOD), Brent Saunders (BOD)

The platform is delivering



Discover
Novel genetic disease targets

20+

Disclosed programs in the pipeline



Create
Medicines with industry-leading research capabilities

>10

INDs since 2015



Test
Our drugs through global development footprint

18

Clinical trials across the globe



Deliver
Our products to patients through commercial infrastructure

2

Product launches expected in 2021


BridgeBio corporate overview



Strategy



Platform



Products



Our pipeline of 20+ development programs spans multiple therapeutic areas and drug modalities

 Small molecule
  Topical small molecule
  Biologics
  Gene therapy

Portfolio segment	Program	Drug mechanism	Diseases	Patient pop. (US+EU)	Modality	Preclinical		Clinical		
						Discovery	IND-enabling	Phase1	Phase 2	Phase 3
Mendelian 	Acoramidis	TTR stabilizer	ATTR-CM	>400K						
	Fosdenopterin	cPMP replacement	MoCD type A	100						
	Infigratinib	Low-dose FGFR1-3i	Achondroplasia	55K						
	Encaleret	CaSR antagonist	ADH1 / HP	12K ¹ / 200K						
	Zuretinol	Synthetic retinoid	IRD (RPE65 or LRAT)	3K						
	BBP-418	Glycosylation substrate	LGMD2i	7K						
	BBP-711	GO1 inhibitor	PH1 / FSF	5K / 1.5M						
	BBP-671	PanK activator	PKAN / OA	7K						
	BBP-761	Succinate prodrug	LHON	20K						
	BBP-472	PI3Kβi	PTEN autism	120K						
Genetic Dermatology 	Patidegib²	Topical SMOi	Gorlin / BCC	120K						
	BBP-589	Recombinant COL7	RDEB	1.5K						
	BBP-681	Topical PI3Kai	VM / LM	117K						
	BBP-561	Topical KLK 5/7i	Netherton	11K						
Targeted Oncology 	Infigratinib	FGFR1-3i	FGFR+ tumors	37K						
	BBP-398	SHP2i	Multiple tumors	>500K						
	BBP-454	Pan-mutant KRASi	KRAS+ tumors	>500K						
	BBP-954	GPX4i	Multiple tumors	>500K						
Gene Therapy 	BBP-631	21-OH gene therapy	CAH	>75K						
	BBP-812	ASPA gene therapy	Canavan	1K						
	BBP-815	TMC1 gene therapy	Genetic hearing loss	10K						

NDA filed

1 US carriers; 2 We are party to an option agreement pursuant to which LEO Pharma A/S has been granted an exclusive, irrevocable option to acquire PellePharm, including the BBP-009 program. If the option is exercised by LEO Pharma A/S, we will no longer have rights to develop and commercialize BBP-009.

Four core value drivers over the next 12-18 months

Program	Patient population	Status	Upcoming event(s)
Acoramidis: TTR stabilizer for ATTR	 >400K	ATTR-CM Ph3 ongoing	☐ Topline Ph3 part A data late-2021 / early-2022 ☐ Topline Ph3 part B data 2023
Low-dose infigratinib (FGFRi) for achondroplasia	 55K	Enrolling Ph2 study	☒ Dose first child ☐ Phase 2 data 2021
Gene therapy for congenital adrenal hyperplasia (BBP-631)	 >75K	GLP tox ongoing	☐ File IND ☐ Phase 1/2 data 2021
Encaleret: CaSR inhibitor for autosomal dominant hypocalcemia type 1 (ADH1)	 12K	Ph2 ongoing	☒ FPI in Ph2 study ☐ Phase 2 data 2021

\$711mn in cash and equivalents as of September 2020 expected to provide runway into 2022

A pipeline with multi-blockbuster potential



\$1B+
opportunities
in the pipeline

- 1) **Acoramidis** for ATTR CM and PN
- 2) **Low-dose infigratinib** for achondroplasia
- 3) **AAV5 gene therapy** for congenital adrenal hyperplasia
- 4) **High-dose infigratinib** for adjuvant urothelial carcinoma
- 5) **Pan-mutant KRAS inhibitor** for KRAS+ cancer
- 6) **SHP2 inhibitor** for RAS and kinase mutant cancer
- 7) **GPX4 inhibitor** for multiple tumor types
- 8) **GO1 inhibitor** for frequent kidney stone formers



Acoramidis (AG10) for transthyretin (TTR) amyloidosis (ATTR)

Seeking to address large and growing need in ATTR, a progressive and fatal disease affecting >400K patients

Designed to target the disease at its source by stabilizing TTR, a genetically and clinically validated mechanism


Advancing acoramidis, a potential best-in-class drug that mimics naturally occurring rescue mutation

Acoramidis has been **well-tolerated and demonstrated near-complete TTR stabilization** in Phase 1 and Phase 2 studies

Executing Phase 3 study with top-line data expected in late-2021 / early-2022

Art
ATTR-CM patient

Acoramidis was designed to treat ATTR at its source



Acoramidis has been well-tolerated and demonstrated near-complete TTR stabilization in pre-clinical, Ph1, and Ph2 studies

Phase 2 safety summary¹


	Placebo N = 17	Acoramidis (pooled doses) N = 32
Any Adverse Event	15 (88%)	21 (66%)
Mild	6 (35%)	11 (34%)
Moderate	8 (47%)	9 (28%)
Severe	1 (6%)	1 (3%)
Any Serious Adverse Event	2 (12%)	1 (3%)
AF and CHF	1 (6%) ¹	0
Leg cellulitis	1 (6%)	0
Dyspnea	0	1 (3%)

¹ Judge, D.P. et al. JACC Vol. 74, No. 3, 2019:285 – 95

² Judge, D.P. et al. American Heart Association 2019


Phase 2 TTR stabilization²

TTR stabilization at steady-state trough level
%, mean ± SEM




Deaths and CV hospitalizations reported in acoramidis Phase 2 OLE were lower than in placebo-treated ATTR-ACT participants

All-cause mortality at 15 months
Participants died or receiving transplant (%)



Cardiovascular hospitalizations at 15 months
Participants with ≥1 CV hospitalization (%)



1 Based on routine adverse event reporting

Note: These data are based on a cross-trial comparison and not a randomized clinical trial. As a result, the values shown may not be directly comparable

Source: Judge, DP et al. American Heart Association Scientific Sessions 2019

ATTRibute-CM will provide 12-month functional outcome data and 30-month mortality and CV hospitalization data

Key inclusion criteria

- Subjects with diagnosed ATTR-CM (WT or mutant)
- NYHA Class I-III
- ATTR-positive biopsy or ^{99m}Tc scan
- Light chain amyloidosis excluded if diagnosis by ^{99m}Tc

Screening and randomization

12-month primary endpoint:
Change in 6MWD

30-month primary endpoint:
Mortality and CV hospitalizations

800 mg acoramidis twice daily

Target N ~ 340

Placebo twice daily

Target N ~170

800 mg
acoramidis
twice daily


Part A

Part B
Tafamidis usage allowed

Open label extension

6MWD = Six-minute walk distance; NYHA = New York Heart Association;
^{99m}Tc = Technetium labeled pyrophosphate (PYP) or bisphosphonate (e.g., DPD);
CV = cardiovascular-related


ATTRibute-CM is designed to evaluate safety and efficacy of acoramidis across complementary measures of drug activity and ATTR-CM disease progression



NT-proBNP = N-terminal pro b-type natriuretic peptide; TnI = Troponin I; 6MWD = Six-minute walk distance
QoL = Quality of life

Rapid functional decline in untreated ATTR-CM patients provides opportunity to demonstrate robust clinical benefit

Summary of 6MWD data in ATTR-CM and healthy cohorts




Optimal profile for tafamidis would markedly slow or halt decline in 6MWD in trial participants

1. Enright, P.L. et al. Chest 2003. N = 3333 healthy elderly adults, baseline set to match ATTR-ACT placebo group
2. Maurer, M.S. et al. NEJM 2018. N = 264 (tafamidis), N = 177 (placebo) ATTR-CM trial participants
3. Lane, T.L. et al. Circulation 2019. N = 1034 ATTR-CM patients

Part B endpoint will hierarchically compare mortality and cardiovascular hospitalizations between all pairs of trial participants

Schematic illustration of win ratio analysis¹



1 Primary analysis of ATTRibute-CM will use a modified win ratio analysis (Finkelstein-Schoenfeld)

Ongoing and planned studies of acoramidis aim to continually expand clinical evidence and addressable patient population

ATTRibute-CM Phase 3 study expected to enroll over 600 participants and on track for topline data in late 2021 or early 2022

2021/
2022

ATTRibute^{CM}

ATTR-CM
WT and hereditary
Functional outcomes

2023

ATTRibute^{CM}

ATTR-CM
WT and hereditary
Functional outcomes
+
Composite mortality and morbidity

2024

ATTRibute^{PN}

ATTR-PN
Hereditary
Functional outcomes

ATTRibute^{PN}

ATTR-PN
Hereditary
Functional outcomes

2025+

Prevention in high risk populations
Head-to-head comparisons



Claudia,
child with achondroplasia

Low-dose FGFR inhibitor (infigratinib) for achondroplasia

Achondroplasia overview



Prevalence

55,000 (US+EU) –
one of the most common
genetic conditions



Genetic driver

FGFR3 activation



Pathophysiology

Up-regulation of STAT1 and MAPK
in the growth plate cause cranial,
spinal, and stature symptoms

Features of a potential best-in-class medicine for achondroplasia



Direct targeting of FGFR3

and normalization of both
STAT1 and MAPK
signaling pathways




Potential to address all

drivers of symptoms,
including cranial, spinal
and stature issues



Oral dosing, the most
convenient solution for
children with achondroplasia
and their families

Potential best-in-class approach targeting achondroplasia directly at its genetic source



ACH FGFR3 gain-of-function mutation causes:

- 2-3x over-activation of the receptor
- Up-regulation of downstream pathways STAT1 and MAPK
- Aberrant growth plate development, which causes cranial, spinal, and stature symptoms

Low-dose infigratinib has the potential to:

- Directly inhibit the causal gain-of-function mutation in FGFR3
- Normalize both the STAT1 and MAPK signaling pathways
- Reverse all key drivers of symptoms

Low-dose infigratinib improves all the key drivers of clinical symptomology in validated ACH mouse model

1 Cranial bone issues

17%

increase in FM area

6%

increase in AP skull length




May lead to **decrease in foramen magnum stenosis** and fewer surgeries

FGFR3 WT
No treatment

FGFR3^{Y367C/+}
No treatment

FGFR3^{Y367C/+}
Infigratinib tx



2 Disorders of the spine

12%


increase in L4-L6 length

73%

increase in disc width



May lead to **decrease in spinal stenosis**, possibly **reducing need for surgery**



3 Disproportionate short stature

21%


increase in femur length

33%

increase in tibia length




May lead to **increased stature and proportionality**



Source: Komla-Ebri et al. J Clin Inv 2016

Note: percent increase compared to vehicle treated FGFR3^{Y367C/+} mouse, infigratinib treatment with 2mg/kg subcutaneous dos

Low-dose infigratinib showed potential best in-class preclinical profile in validated achondroplasia mouse model

Company/ Asset	MOA	Route	Status	Mouse model	Tibia length	Femur	Foramen magnum area	L4-L6 height
 Infigratinib	Selective FGFR1-3i	Oral	Ph2	FGFR3 ^{Y367C}	 32.6%	 20.9%	 17.0%	 12.1%
 Vosoritide (BMN111)	CNP analogue	Daily SQ	Pivotal (NDA filed)	FGFR3 ^{Y367C/+}	 6.6%	 5.2%	No known publicly available data	 3.3%
 TransCon CNP ¹	CNP analogue	Weekly SQ	Ph2	FGFR3 ^{Y367C/+}	 12.3%			
 Reifercept (TA-46)	FGFR3 decoy	Weekly SQ	Ph1	FGFR3 ^{ACH}	 8.6%	 6.2%		

Preclinical data from infigratinib and other investigational achondroplasia therapies


Percent increase compared to non-treated mouse

Source: Komla-Ebri et al. J Clin Inv 2016, Lorget et al. Am J Hum Genet 2012, Garcia et al. Science Trans Med 2013, Breinholt ENDO 2017
Note: subcutaneous doses, percent increase compared to vehicle treated FGFR3^{Y367C/+}, FGFR3^{ACH/+} mouse as noted in “Mouse model” columns
Infigratinib treatment with 2mg/kg subcutaneous dose ¹Based on vosoritide continuous infusion; *Value estimated using Digitizelt.

We have a wide anticipated therapeutic index in achondroplasia


Infigratinib has been tested in >700 humans in our oncology program, providing significant data on PK, tolerability and safety

Most common and dose-limiting side effect is phosphorus elevation (on-target through FGFR1 inhibition), which occurs significantly above our planned achondroplasia doses



¹Based on 125mg dose and 60kg adult; ²Based on estimated TD₅₀ at 40mg and 60kg adult; ³Based on PK modeling and allometric scaling from animal models

The PROPEL clinical program is enrolling with data expected in 2021



Key inclusion criteria

- Children 2.5 – 10 years old
- Clinical and molecular ACH diagnosis

Primary objectives

- Baseline annualized growth velocity (AGV)

Primary objectives

- Identify safe therapeutic dose for expansion / pivotal study
- Safety and tolerability
- Change from baseline in AGV

Primary objectives

- Long-term safety and efficacy

Pipeline



BBP-631: AAV5 for CAH
(IND-enabling)
IND anticipated in 2020



BBP-812: AAV9 for Canavan
Disease
IND anticipated in 2020



BBP-815: AAV for TMC1
hearing loss



Multiple undisclosed
discovery programs

BridgeBio's gene therapy programs and capabilities



Collaborations with leading gene therapy pioneers



In-house research vector manufacturing and optimization



Best-in-class CMC, process development, and analytical development team



Dedicated clinical/commercial manufacturing space at Catalent



Leading clinical, non-clinical and regulatory team, with significant gene therapy experience

BBP-631: AAV5 gene therapy for congenital adrenal hyperplasia (CAH)

Program overview



Prevalence

75,000 (US+EU) – One of the largest known AAV gene therapy markets



Genetic driver

21-hydroxylase inactivation



Pathophysiology

Inability to produce cortisol causes need for supraphysiologic doses of synthetic steroids, 3x increase in mortality risk, hirsutism, Cushingoid symptoms


We believe CAH is an ideal indication for AAV gene therapy:

- **Low threshold to correct phenotype**, validated by human clinical genetics (~5-10% of WT enzyme activity)
- **Only approach designed to induce endogenous cortisol and mineralocorticoid production**, potentially allowing steroid withdrawal
- **Durable transgene delivery to the adrenal gland of NHPs** with IV dosing of our construct
- **Preliminary Ph1/2 data anticipated in 2021** with endogenous cortisol production as a key endpoint

Maris,
child with CAH


Gene therapy is the only modality designed to treat CAH at its source and allow for production of endogenous cortisol

Healthy Hypothalamic-Pituitary-Adrenal Axis



In a functional HPA system, cortisol and aldosterone are produced as needed by the body. Cortisol serves as a “brake” on the CRF/ACTH system

Hormonal dysregulation with 21OHD; no cortisol “brake” on ACTH, shunting of 17OHP to androgens




In CAH, cortisol and aldosterone are not able to be produced. The lack of a “cortisol brake” results in buildup of progesterone and 17OHP, leading to an excess of androgen production

CAH patients have 3-4X higher mortality than the general population, and suffer significant morbidity ranging across cardiovascular and metabolic disease, bone disease, infertility, chronic fatigue, and other disorders.


CAH: NHP study showed durable transgene expression; 5-10% of WT enzyme may be sufficient for clinical impact

Genotype-phenotype studies show that >5-10% of enzyme activity results in nonclassical CAH




- Due to the high enzymatic efficiency/selectivity of 21-OHase, **only a small amount of enzyme is required to rescue the phenotype**

Mouse studies show a VGC of only 0.13 at 18 wks was sufficient for phenotypic correction



- At 15 weeks in treated mice, **progesterone** (the key substrate of 21OHase in mice) was **significantly reduced vs untreated mice**

NHP studies show sustained VGC and RNA out to 6 months




- Transgene expression is dose-dependent and stable out at 24 wks
- We can durably transduce the NHP adrenal gland with our construct at >20x the vector required to correct the CAH phenotype in mice

NHP protein data using mass spec methods suggests potentially therapeutic levels of 21-hydroxylase enzyme

Human 21-hydroxylase protein as a % of NHP 21-hydroxylase protein (Mass Spec quantification)

- We have developed mass-spec methods to quantify protein expression by identifying differential peptides between human and NHP 21-OH
- These data suggest dose-dependent enzyme expression in the adrenal cortex from 9%-24% of WT levels
- Genotype-phenotype relationship suggests as little as 5% of WT enzyme activity is associated with the mild/asymptomatic non-classic form of CAH





Encaleret for disorders of calcium homeostasis, including autosomal dominant hypocalcemia type 1 (ADH1)

Targets hypocalcemia/hypercalciuria by selectively antagonizing the calcium-sensing receptor (CaSR)

- Opportunity identified in collaboration with experts at the NIH

Prior clinical experience enables accelerated development


- Well tolerated in >1,200 human subjects excepting dose-dependent increases in serum calcium (target effect in ADH1 patients)
- Phase 2 study in ADH1 planned to initiate in 2020 with proof-of-concept data anticipated in 2021

Potential 1st in class CaSR antagonist with differentiated profile for ADH1 and hypoparathyroidism

- Initial development in genetically-defined population of ADH1, driven by CaSR activating mutations (~12K carriers in US)
- Potential for expansion into post-surgical chronic hypoparathyroidism (~200K patients in US & EU)

**Alexis and Jackson
ADH1 patients**

Encaleret is designed to treat ADH1 at its source



The calcium sensing receptor (CaSR) primarily acts to regulate parathyroid hormone levels and renal calcium reabsorption

Illustration of parathyroid cell





Illustration of renal tubule





Encaleret is designed to target ADH1 at its source by normalizing hyper-active calcium sensing receptor


Rationale for studying CaSR inhibitor in ADH1

- ADH1 is caused by activating mutations in the CaSR leading to hypocalcemia and hypercalciuria
- Prior generation CaSR inhibitor partially addressed ADH1 phenotype despite limited exposure¹


Encaleret normalized serum and urine calcium in a mouse model of ADH1²



Encaleret increased serum calcium in clinical trials in patients with osteoporosis³



Phase 2, open-label dose-ranging study will evaluate safety, tolerability, and efficacy of encaleret in ADH1



Top-line, proof-of-concept results of encaleret in ADH1 are anticipated in 2021

¹ Pharmacodynamic measurements to be collected through duration of study



BridgeBio oncology research

World-class oncology team drives our discovery and development

Eli Wallace

CSO Oncology Research



Pedro Beltran

SVP Oncology



Frank McCormick

Chairman of Oncology



Richard Scheller


Chairman of R&D



Basia, pancreatic cancer
patient (>90% KRAS-driven)


Three disclosed oncology research targets

SHP2 (BBP-398)




- Potential best in class oral compound
 - Optimized safety, PK and PD profile
 - Maximizes combination therapy potential
- IND cleared

KRAS



- Multiple unexploited sites
- Comprehensive pan-mutant targeting approaches

GPX4



- Potential first in class compound for novel cancer target
- In vivo monotherapy activity and combo potential

Partnerships afford us exceptional collaborators and resources




- Partnership with the National RAS Initiative, including 60 of the world's **foremost academic RAS researchers**
- Cutting edge RAS **structural biology expertise**
- Utilization of **cutting-edge instrumentation and techniques**, as well as the **expertise** to lead experiments



- Home to Sierra: the **world's 3rd fastest computing system**
- Enables **multi-microsecond molecular dynamics simulations** of protein complexes, and highly efficient *in silico* **docking simulations**
- This computing power, combined with RAS structural biology expertise at the NCI, delivers **unique insights that fuel our drug design**

Crystal structure enables a static understanding of the target ...

KRAS4b model based on crystal



G-domain
G-domain switch I


G-domain switch II
Hypervariable region

One therapeutic approach is to inhibit KRAS4b **membrane localization** by targeting **hypervariable region**

Static model reveals only a **subset of potential binding sites** for pharmacological compounds

... whereas molecular dynamics simulation reveals transient conformations and interactions

KRAS4b simulation



G-domain
G-domain switch I

G-domain switch II
Hypervariable region

Reveals possible KRAS4b **HVR transient localization to G-domain**

Elucidates potential transient druggable pocket where **compounds could react covalently with C185**

Enables *in silico* SAR to **inhibit KRAS4b membrane localization**


KRAS: multiple shots on goal with our pan-mutant inhibitor programs – each with a unique MOA targeting a novel pocket

KRAS pathway in cancer	Program	MOA	Targets KRAS GTP	Pan-mutant	Crystal structure	Molecular Dynamics
Receptor tyrosine kinase signals (EGFR, ALK, TRK, RET) Cancer mutations lock KRAS in its active state SHP2 Prenylation C185 KRAS GTP RAS(ON) H95 RAS effectors RAF PI3K Tumor cell proliferation and survival	Program 1: H95 targeting	- Directly binds activated KRAS through H95 - Inhibits KRAS from signaling through effectors	✓	✓	✓	✓
	Program 2: PI3K effector blocking	- Blocks specific interaction between KRAS and PI3Ka - Blocks PI3K / AKT effector signaling	✓	✓	✓	✓
	Program 3: C185 targeting	- Blocks KRAS from tethering - Blocks conversion of inactive KRAS GDP to active KRAS GTP	✓	✓		✓

Our programs are designed to address all KRAS driver mutations, which occur in >30% of all cancers

SHP2: Our compound shows best-in-class potential and is entering the clinic in 2020

Our SHP2i blocks downstream MAPK signaling and abrogates T cell exhaustion



We believe BBP-398 has the ideal properties for combination with a multitude of other therapeutic classes

Human half life: 10 hours¹

- Allows for recovery above EC50 and reduced MAPK-driven tox

SHP2i combination potential

US + EU incidence, '000s

Potentially differentiated safety profile for combination therapy


- hERG IC50 (μM)*: >100: No QT prolongation

Supporting evidence²

Peclin data:

BBIO SHP2i

Other SHP2i




~1 million patients annually

¹ Predicted human PK based on preclinical in vivo data ² Preclinical data of combination efficacy with SHP2i

SOURCE: US incidence estimated from SEER, TCGA and Kiuru & Busam "The NF1 gene in tumor syndromes and melanoma"; all scaled for WW incidence

SHP2: BBP-398 monotherapy study initiating in 2020; combo trials to follow



Multiple catalysts anticipated through year-end 2021

			ESTIMATED
2020		2021	
1H	2H	FY	
✓ New program announcements ✓ FGFRi for cancer: FPI Ph3 adjuvant urothelial carcinoma study ✓ FGFRi for cancer: FPI Ph2 FGFR fusion tumor agnostic Ph2 study	✓ Low-dose FGFRi for achondroplasia: Begin dosing Ph2 ✓ New IND filings ✓ cPMP for MoCD type A: Complete NDA submission ✓ CaSR antagonist for ADH1: FPI in Ph2 study □ FGFRi for cancer: 2L CCA NDA filing	□ TTR stabilizer for ATTR: Ph3 part A data late-2021 / early-2022 □ Low-dose FGFRi for achondroplasia: Ph2 PoC data □ CAH gene therapy: Ph1/2 PoC data □ CaSR antagonist for ADH1: Ph2 POC data □ Topical SMOi for Gorlin: Topline Ph3 data □ FGFRi for cancer: 2L CCA approval and launch □ cPMP for MoCD type A: Approval and launch □ Recombinant COL7 for RDEB: Topline Ph1/2 data	

\$711mn in cash and equivalents as of September 2020 expected to provide runway into 2022